

QUANTUM DEFORMATIONS OF THE SELF-DUALITY EQUATION  
AND CONFORMAL TWISTORS

*B.M.Zupnik*

*Bogoliubov Laboratory of Theoretical Physics , JINR, Dubna, Moscow Region, 141980,  
Russia, E-mail: zupnik@thsun1.jinr.dubna.su*

Submitted to Proceedings of VI seminar "Quantum Gravity", Moscow, June 12-19, 1995

**Abstract.** A noncommutative algebra of the complex  $q$ -twistors and their differentials is considered on the basis of the quantum  $GL_q(4) \times SL_q(2)$  group. Real and pseudoreal  $q$ -twistors are discussed too. We consider the quantum-group self-duality equation in the framework of the local gauge algebra of differential forms on  $q$ -twistor spaces. Quantum deformations of the general multi-instanton solutions are constructed. The corresponding noncommutative algebras of moduli are introduced. The general  $q$ -instanton connection is a function of the  $q$ -twistors and the  $q$ -moduli .

# 1 Introduction

Noncommutative generalizations of the local gauge theories have been considered in the framework of different approaches [1-7]. The mathematically strict approach of Refs[3,4] is based on the noncommutative global generalizations of the classical fibre bundles. We prefer to study the local structure of the quantum-group gauge theory in terms of the deformed connection and curvature differential forms  $A$  and  $F$ . The basic algebra of these gauge forms should be covariant under the action of the quantum gauge group [5,6]

$$A \rightarrow TAT^{-1} + dTT^{-1} \quad (1.1)$$

where  $T$  and  $dT$  are elements of the differential complex on the quantum group.

Consider some classical or quantum space with the coordinates  $z$  and let  $T(z)$  and  $dT(z, dz)$  be noncommutative 'functions' on this space. We shall treat these functions as generators of the local gauge differential complex if the map

$$T \rightarrow T(z), \quad dT \rightarrow dT(z, dz) \quad (1.2)$$

conserves all relations between  $T$  and  $dT$ .

The general function  $T(z)$  is some formal expansion with noncommutative coefficients. Thus, the localization of the quantum group is equivalent to the definition of an infinite-dimensional noncommutative Hopf algebra.

The  $q$ -deformations of the Grassmann and twistor spaces were studied in Refs[8-10]. In section 2 we consider the differential calculus on the 4-dimensional deformed complex twistor space  $T_q(4, C)$ . The real forms of the  $q$ -twistor space are also discussed.

Section 3 is devoted to the description of the quantum-group gauge fields on the  $q$ -twistor space. We consider the algebraic relation for the connection form in the  $GL_q(N)$  gauge theory [5,6] that defines the algebraic properties of the 'off-shell' gauge fields. The generalizations of the Yang-Mills and self-duality equations are discussed. Note that one can use the 'pure gauge'  $U(1)$  field and  $q$ -traceless curvature 2-forms in the  $U_q(N)$  gauge theory [7]. The noncommutative analogue of the BPST one-instanton solution [11] was constructed in the deformed 4-dimensional Euclidean space [7].

The deformed analogue of the t'Hooft multi-instanton twistor solution for the gauge group  $GL_q(2)$  (or  $U_q(2)$ ) is considered in section 4. We use a multidimensional extension of the  $q$ -twistor algebra by the set of noncommutative  $6D$ -vector generators  $b$ . The potential of our solution is a sum of the central  $(z, b)$ -functions obeying the  $q$ -twistor Laplace equation.

The deformed generalization of the Atiyah-Drinfel'd-Hitchin-Manin solution [12] for the gauge group  $GL_q(N)$  contains  $q$ -twistor functions  $u$  and  $\tilde{u}$ . We generalize the classical conformal constructions of Ref[13]. One can consider the linear twistor functions  $v$  and  $\tilde{v}$  that depend on the noncommutative moduli  $b$  and  $\tilde{b}$ . The functions  $u$  and  $v$  are submatrices of the quantum  $GL_q(N + 2p)$  matrix for the instanton number  $p$ . The consistency relations for the deformed ADHM-construction can be proved in the framework of a differential calculus on  $GL_q(N + 2p)$ . The self-duality condition is equivalent to the bilinear constraint on the moduli  $b$  and  $\tilde{b}$ .

A preliminary version of this work was published in Ref[14]. Note that we use here the modified notation and definitions of some basic quantities.

## 2 Differential calculus on the deformed twistor space

The conformal covariant description of the classical ADHM solution was considered in Ref[13]. This approach uses real forms of the complex  $GL(4, C) \times SL(2, C)$  twistors where  $GL(4, C)$  is the complex conformal group. It is convenient to discuss firstly the deformed complex twistors.

Let  $R_{cd}^{ab}$  be a solution of the Yang-Baxter equation satisfying also the Hecke relation

$$R R' R = R' R R' \quad (2.1)$$

$$R^2 = I + (q - q^{-1})R \quad (2.2)$$

where  $q$  is a complex parameter. Note that the standard notation for these  $R$ -matrices is  $R = \hat{R}_{12}$ ,  $R' = \hat{R}_{23}$  [15]. We use the symbols  $a, b \dots h = 1 \dots 4$  for the 4D-spinor indices.

The multiparameter 4D  $R$ -matrix [16-18] and corresponding inverse matrix  $R^{-1}$  can be written in the following simple form:

$$(R^{\pm 1})_{cd}^{ab} = \delta_c^a \delta_d^b [q^{\pm 1} - q^{\epsilon(a-b)}] + r(ab) \delta_d^a \delta_c^b \quad (2.3)$$

where  $\epsilon(b-a) = 0$ ,  $\pm 1$  is a sign function and  $r(ab)$  are complex parameters satisfying the relations  $r(ab)r(ba) = 1$ . It is evident that this formula is valid for an arbitrary number  $N$ .

The standard  $GL_q(4)$  solution corresponds to the case  $r(ab) = 1$  [15]. The choice  $q = 1$  leads to the unitary  $R$ -matrix [19]

$$R^2 = I \quad (2.4)$$

Consider also the  $SL_q(2, C)$   $R$ -matrix

$$R_{\mu\nu}^{\alpha\beta} = q \delta_\mu^\alpha \delta_\nu^\beta + \varepsilon^{\alpha\beta}(q) \varepsilon_{\mu\nu}(q) \quad (2.5)$$

where  $\varepsilon(q)$  is the deformed antisymmetric symbol

$$\varepsilon^{12}(q) = -\varepsilon_{12}(q) = \frac{1}{\sqrt{q}}, \quad \varepsilon^{21}(q) = -\varepsilon_{21}(q) = -\sqrt{q} \quad (2.6)$$

The  $q$ -deformed flag spaces and twistors were considered in Refs[8-10]. We shall treat the complex  $q$ -twistors  $z_a^\alpha$  as generators of the noncommutative algebra with the basic relation

$$R_{\mu\nu}^{\alpha\beta} z_a^\mu z_b^\nu = z_c^\alpha z_d^\beta R_{ba}^{dc} \quad (2.7)$$

This relation for the  $(4 \times 2)$  rectangular matrix  $z$  is analogous to the RTT-relations for the square quantum matrices. The consistency conditions for (2.7) are pairs of the Yang-Baxter and Hecke relations (2.1, 2.2) for the independent 4D and 2D  $R$ -matrices with the unique common parameter  $q$ .

The differential calculus on the complex  $q$ -twistor space  $T_q(4, C)$  can be constructed by the analogy with the bicovariant differential complex on the quantum linear group [20-23]. Consider the relations between  $z_a^\alpha$  and their differentials  $dz_a^\alpha$

$$z_a^\alpha dz_b^\beta = R_{\mu\nu}^{\alpha\beta} dz_c^\mu z_d^\nu R_{ba}^{dc} \quad (2.8)$$

$$dz_a^\alpha dz_b^\beta = -R_{\mu\nu}^{\alpha\beta} dz_c^\mu dz_d^\nu R_{ba}^{dc} \quad (2.9)$$

The elements  $z$  and  $dz$  are generators of the external algebra  $\Lambda T_q(4, C)$  on the  $q$ -twistor space. The operator of external derivative  $d$  on  $\Lambda T_q(4, C)$  is nilpotent and satisfies the ordinary Leibniz rule.

The symmetry properties of  $dz$  can be obtained from Eq(2.9)

$$P_2^{(+)} dz dz' P_4^{(+)} = 0 = P_2^{(-)} dz dz' P_4^{(-)} \quad (2.10)$$

where  $P_2^{(\pm)}$  and  $P_4^{(\pm)}$  are the projection operators for  $SL_q(2)$  and  $GL_q(4)$ , respectively [15]

$$P^{(+)} + P^{(-)} = I, \quad R = qP^{(+)} - q^{-1}P^{(-)} \quad (2.11)$$

One can define the algebra of partial derivatives  $\partial_\alpha^a$  on  $T_q(4, C)$

$$R_{cd}^{ab} \partial_\alpha^c \partial_\beta^d = \partial_\mu^a \partial_\nu^b R_{\beta\alpha}^{\nu\mu} \quad (2.12)$$

$$\partial_\alpha^a z_b^\beta = \delta_b^a \delta_\alpha^\beta + R_{\alpha\nu}^{\beta\mu} R_{cb}^{da} z_d^\nu \partial_\mu^c \quad (2.13)$$

$$\partial_\alpha^a dz_b^\beta = R_{\alpha\nu}^{\beta\mu} R_{cb}^{da} dz_d^\nu \partial_\mu^c \quad (2.14)$$

Consider a definition of the deformed  $\varepsilon_q$ -symbol for  $GL_q(4)$

$$\varepsilon_q^{abcd} = -q R_{fe}^{ba} \varepsilon_q^{efcd} = [P_4^{(-)}]_{fe}^{ba} \varepsilon_q^{efcd} = -q^{\epsilon(b-a)} r(ba) \varepsilon_q^{bacd} \quad (2.15)$$

Analogous relations are valid for other neighboring pairs of indices.

The  $q$ -twistors obey the following identity:

$$\varepsilon_q^{abcd} z_b^\beta z_c^\mu z_d^\nu = 0 \quad (2.16)$$

Introduce the  $SL_q(2)$ -invariant bilinear function of  $q$ -twistors

$$y_{ab} = \frac{q^2}{1+q^2} \varepsilon_{\alpha\beta}(q) z_a^\alpha z_b^\beta = [P_4^{(-)}]_{ba}^{dc} y_{cd} \quad (2.17)$$

This vector satisfies the following commutation relation

$$y_{ab} z_c^\alpha = q^{-1} R_{ha}^{ed} R_{cb}^{fh} z_d^\alpha y_{ef} \quad (2.18)$$

In consequence of Eq(2.16) the coordinate  $y$  is an isotropic vector in the deformed 6D space

$$(y, y) = \varepsilon_q^{abcd} y_{ab} y_{cd} = 0 \quad (2.19)$$

This  $GL_q(4, C)$  covariant equation determines the 4D deformed subspace of the complex 6D quantum plane. The classical analogue of this subspace is a complex sphere  $S_4^C$ .

Consider a duality transformation  $*$  of the basic  $q$ -twistor 2-forms by analogy with Ref[7]

$$* dz dz' = dz dz' P_4^{(+)} - dz dz' P_4^{(-)} \quad (2.20)$$

where  $P_4^{(\pm)}$  are the  $GL_q(4)$  projectors (2.11). Note that a self-dual part  $dz dz' P_4^{(+)}$  is proportional to the  $SL_q(2)$ -invariant conformal tensor

$$\varepsilon_{\alpha\beta}(q) dz_a^\alpha dz_b^\beta \quad (2.21)$$

Real  $q$ -twistors can be treated as a representation of the real quantum group  $SL_q(2, R) \times GL_q(4, R)$ . The classical analogue of these twistors are connected with the real pseudo-Euclidean  $(2, 2)$ -space [13]. Consider the  $R$ -matrices (2.3, 2.5) and the conditions  $|q| = 1$  and  $|r(ab)| = 1$ , then under the complex conjugation

$$\overline{R_{cd}^{ab}} = (R^{-1})_{dc}^{ba} \quad (2.22)$$

$$\overline{R_{\mu\nu}^{\alpha\beta}} = (R^{-1})_{\nu\mu}^{\beta\alpha} \quad (2.23)$$

These formulas correspond to anti-involution of the real  $T_q(2, 2)$  twistors

$$\overline{z} = z, \quad \overline{dz} = dz \quad (2.24)$$

$$\overline{z z'} = z' z, \quad \overline{z dz'} = dz' z \quad (2.25)$$

The pseudoreal Euclidean  $q$ -twistors have a more complicated anti-involution

$$\overline{z_a^\alpha} = \varepsilon_{\alpha\beta}(q) z_b^\beta C_a^b(q) \quad (2.26)$$

where  $C(q)$  is the charge-conjugation matrix for the Euclidean conformal quantum group  $U_q^*(4) = D \times SU_q^*(4)$  and  $D$  is a real one-parameter dilatation.

It is convenient to use the simple representation

$$C(q) = \begin{pmatrix} \varepsilon^{\alpha\beta}(q) & 0 \\ 0 & \varepsilon^{\dot{\alpha}\dot{\beta}}(q) \end{pmatrix} \quad (2.27)$$

For the real  $q$  we have the following properties:

$$\overline{\varepsilon^{\alpha\beta}(q)} = \varepsilon^{\alpha\beta}(q) = -\varepsilon_{\alpha\beta}(q) \quad (2.28)$$

$$\overline{C(q)} = C(q), \quad C^2(q) = -I \quad (2.29)$$

$$\overline{\overline{z_a^\alpha}} = -\varepsilon^{\alpha\beta}(q) \varepsilon_{\beta\gamma}(q) z_b^\gamma (C^2)_a^b(q) = z_a^\alpha \quad (2.30)$$

$$\overline{R_{ba}^{dc}} = C_e^d(q) C_f^c(q) R_{gh}^{fe} C_a^g(q) C_b^h(q) \quad (2.31)$$

We do not here consider the conformal quantum group  $SU_q(2, 2)$  [24, 25] and corresponding  $q$ -twistors.

### 3 Quantum-group gauge theory on the $q$ -twistor space

The classical gauge field on some domain  $\{x^m\}$  of the basic space corresponds to the connection 1-form

$$A(x, dx) = dx^m A_m(x) \quad (3.1)$$

which can be decomposed in terms of the gauge-group generators. For the domain with the coordinates  $\tilde{x}(x)$  one should define the transformed connection

$$\tilde{A}(\tilde{x}, d\tilde{x}) = T(x)AT^{-1}(x) + dT(x)T^{-1}(x) \quad (3.2)$$

where  $T(x)$  is a matrix of the local gauge transformation.

The components of the matrix  $A(x, dx)$  satisfy the anticommutativity conditions

$$\{A_k^i, A_m^l\} = 0 \quad (3.3)$$

The classical gauge group formally has an infinite number of generators. A constructive example of the classical gauge algebra is the affine (Kac-Moody) algebra. The quantum affine algebras can be considered as a basis of the quantum gauge theory on the classical two-dimensional space.

The formal quantization of the gauge groups on the multi-dimensional classical or quantum spaces is a difficult problem. Let  $R_N$  be the constant  $R$ -matrix for the quantum group  $GL_q(N)$  and  $x^M$  are the coordinates of some basic space. Consider the simplest possible relations for the components of the quantum gauge matrix

$$R_N T(x) T'(x) = T(x) T'(x) R_N \quad (3.4)$$

$$T_k^i(x) = \sum_{n=0}^{\infty} \frac{1}{n!} (T_k^i)_{M_1 \dots M_n} x^{M_1} \dots x^{M_n} \quad (3.5)$$

where  $i, k \dots = 1 \dots N$ .

The quantum-group gauge matrix is well defined if the relation (3.4) generates the consistent set of relations between the coefficients  $(T_k^i)_{M_1 \dots M_n}$ .

Quantum deformations of the  $GL(N)$  gauge connection can be treated in terms of the noncommutative gauge algebra for the components of the deformed connection 1-form [5,6]

$$(AR_N A + R_N AR_N AR_N)_{mn}^{ik} = A_l^i (R_N)_{rn}^{lk} A_m^r + (R_N)_{jl}^{ik} A_r^j (R_N)_{st}^{rl} A_p^t (R_N)_{mn}^{pt} = 0 \quad (3.6)$$

These relations generalize the classical anticommutativity conditions (3.3). The gauge algebra is an analogue of the relations between components of the right-invariant 1-forms  $\omega = dTT^{-1}$  in the framework of the bicovariant differential calculus on  $GL_q(N)$  [20-23]

$$\omega R_N \omega + R_N \omega R_N \omega R_N = 0, \quad d\omega - \omega^2 = 0 \quad (3.7)$$

Thus, the form  $\omega$  can be considered as a pure gauge  $GL_q(N)$ -field. The general  $GL_q(N)$  connection  $A$  has the nontrivial curvature 2-form

$$F = dA - A^2 \quad (3.8)$$

Explicit constructions of the deformed gauge fields on the  $q$ -twistor space contain also the noncommutative elements (moduli) which generate some algebra  $B$

$$A_k^i(z, dz, B) = dz_a^\alpha (A_\alpha^a)_k^i(z, B) \quad (3.9)$$

The appearance of additional noncommutative elements is necessary for the consistency of the algebra (3.6) and the relations of the  $q$ -twistor algebra (2.8, 2.9).

The (anti)self-duality equation for the gauge field (3.9) can be defined with the help of the relations (2.20)

$$*F = \frac{1}{2}(*dz_a^\alpha dz_b^\beta) F_{\alpha\beta}^{ab}(z, B) = \pm F \quad (3.10)$$

or in terms of the deformed field-strength

$$[P_4^{(\pm)}]_{cd}^{ab} F_{\alpha\beta}^{cd}(z, B) = 0 \quad (3.11)$$

The solutions of this equation and the explicit construction of the algebra  $B$  will be considered in sections 4 and 5.

A quantum deformation of the Yang-Mills equation has the standard form in the framework of the external algebra  $\Lambda T_q(4, C)$

$$\nabla * F = d * F + [A, *F] = 0 \quad (3.12)$$

The bicovariant differential calculus with the ordinary Leibniz rule for the operator  $d$  and the gauge-connection algebra (3.6) are consistent only for the case of the nonsemisimple quantum group  $GL_q(N)$ . The gauge algebra produce the restriction

$$\alpha = \text{Tr}_q A \neq 0 \quad (3.13)$$

Nevertheless, one should use the gauge-covariant conditions [7]

$$d\alpha = 0, \quad \text{Tr}_q A^2 = 0, \quad \text{Tr}_q F = 0 \quad (3.14)$$

These restrictions generate the effective reduction of the Abelian gauge field  $\alpha$  in the framework of the gauge group  $GL_q(N)$ .

## 4 Quantum deformations of the t'Hooft multi-instanton solution

A simple form of the manifest multi-instanton solution in the Euclidean space was discussed in Refs[26-28]. This solution can be written in terms of the potential  $\Phi$  satisfying the Laplace equation. The classical twistor version of the t'Hooft solution was considered in Ref[13].

Consider firstly the deformed Laplace equation in the complex  $q$ -twistor space  $T_q(4, C)$ . Using Eq(2.13) one can obtain the action of the partial  $q$ -twistor derivative on the isotropic  $6D$ -vector

$$\partial_\alpha^c y_{ab} = \varepsilon_{\alpha\beta}(q) [P_4^{(-)}]_{ba}^{dc} z_d^\beta \quad (4.1)$$

Introduce the formal differential operator that acts only on the  $6D$  vector variables

$$\partial^{dc} \triangleright y_{ab} = [P_4^{(-)}]_{ba}^{dc} \quad (4.2)$$

Now we can write the following relations:

$$dy_{ab} = \varepsilon_{\alpha\beta}(q) [P_4^{(-)}]_{ba}^{dc} dz_c^\alpha z_d^\beta \quad (4.3)$$

$$d\Phi(y) = dz_c^\alpha \partial_\alpha^c \Phi(y) = dy_{ab} \partial^{ba} \Phi(y) \quad (4.4)$$

$$\partial_\alpha^c \Phi(y) = \varepsilon_{\alpha\beta}(q) z_b^\beta \partial^{bc} \Phi(y) \quad (4.5)$$

The  $SL_q(2, C)$ -invariant analogue of the Laplace operator has the following form:

$$\Delta^{ba} = -\frac{q}{1+q^2} \varepsilon^{\alpha\beta}(q) \partial_\beta^b \partial_\alpha^a \quad (4.6)$$

By definition, we have

$$\Delta^{ba} \triangleright y_{cd} = [P_4^{(-)}]_{dc}^{ba} = \partial^{ba} \triangleright y_{cd} \quad (4.7)$$

In this section we shall use the standard  $GL_q(4, C)$   $R$ -matrix corresponding to Eq(2.3) with  $r(ab) = 1$ . This  $R$ -matrix satisfies the following identity:

$$\varepsilon_q^{abcd} R_{ea}^{a'h} R_{fb}^{b'e} R_{gc}^{c'f} R_{hd}^{d'g} = q \delta_{h'}^h \varepsilon_q^{a'b'c'd'} \quad (4.8)$$

Introduce the additional noncommutative moduli  $b_{ab}^p$  where  $p$  is an arbitrary number

$$(b^p, b^p) = \varepsilon_q^{abcd} b_{ab}^p b_{cd}^p = 0 \quad (4.9)$$

$$y_{ab} b_{cd}^p = R_{ga}^{ea'} R_{cb}^{fg} R_{he}^{c'b'} R_{df}^{d'h} b_{a'b'}^p y_{c'd'} \quad (4.10)$$

$$b_{ab}^p b_{cd}^{\hat{p}} = q^{-2} R_{ga}^{ea'} R_{cb}^{fg} R_{he}^{c'b'} R_{df}^{d'h} b_{a'b'}^{\hat{p}} b_{c'd'}^p \quad (4.11)$$

and  $p \leq \hat{p}$  in the last equation.

This  $(B, y)$ -algebra has the following central elements

$$X_p = (y, b^p) = \varepsilon_q^{abcd} y_{ab} b_{cd}^p \quad (4.12)$$

The commutativity of  $X_p$  with  $y$  and  $b^{\hat{p}}$  can be proved with the help of Eq(4.8).

Let us introduce the commutation relations between  $b^p$ , and  $z$

$$b_{ab}^p z_c^\gamma = R_{ga}^{eh} R_{cb}^{fg} z_h^\gamma b_{ef}^p \quad (4.13)$$

An analogous relation for  $b^p$ , and  $dz$  can be obtained as the external derivative  $d$  of this formula by using  $[d, b^p] = 0$ .

Equation(2.9) generates the relation for  $y$  and  $dz$

$$y_{ab} dz_c^\gamma = q R_{ga}^{eh} R_{cb}^{fg} dz_h^\gamma y_{ef} \quad (4.14)$$

Write the corresponding relation for the elements  $X_p$

$$X_p dz_a^\alpha = q^2 dz_a^\alpha X_p \quad (4.15)$$



Now one can determine the derivative of the central functions

$$\partial_\alpha^a \frac{1}{X_p} = -\frac{1}{q^2 X_p^2} \partial_\alpha^a X_p \quad (4.16)$$

$$\partial_\alpha^a \frac{1}{X_p^2} = -\frac{1+q^2}{q^4} \frac{1}{X_p^3} \partial_\alpha^a X_p \quad (4.17)$$

It is easy to check the following identity for the isotropic vectors  $b^p$  :

$$\varepsilon^{\alpha\beta}(q) \partial_\beta^b X_p \partial_\alpha^a X_p = -q^{-1} \varepsilon^{abcd} b_{cd}^p X_p \quad (4.18)$$

We can obtain the solutions of the deformed Laplace equation ( $q$ -harmonic functions )

$$\Delta^{ba} \frac{1}{X_p} = \frac{1}{q^5(1+q^2)} \frac{1}{X_p^2} \varepsilon^{\alpha\beta}(q) [\partial_\beta^b \partial_\alpha^a X_p - (1+q^2) \frac{1}{X_p} \partial_\beta^b X_p \partial_\alpha^a X_p] = 0 \quad (4.19)$$

By analogy with Ref[13] one can consider the deformed t'Hooft Ansatz for the  $GL_q(2)$  self-dual gauge field

$$A_\beta^\alpha = q^{-3} dz_a^\alpha (\partial_\mu^\alpha \Phi) \Phi^{-1} \varepsilon^{\sigma\mu}(q) \varepsilon_{\sigma\beta}(q) \quad (4.20)$$

$$\text{Tr}_q A = -q^3 d\Phi \Phi^{-1}, \quad \text{Tr}_q dA = 0 \quad (4.21)$$

where the potential function  $\Phi$  for the instanton number  $P$  is a sum of  $q$ -harmonic functions with different elements  $b^p$

$$\Phi = \sum_{p=1}^P \frac{1}{X_p} = \sum_{p=1}^P (y, b^p)^{-1} \quad (4.22)$$

The anti-self-dual part of the corresponding curvature form vanishes in consequence of Eq(4.19)

$$(F - *F)_\beta^\alpha \sim dz_e^\alpha dz_f^\gamma [P_4^{(-)}]_{ac}^{fe} \Delta^{ac} \Phi \Phi^{-1} \varepsilon_{\beta\gamma}(q) = 0 \quad (4.23)$$

Note that the isotropic vector  $b^p$  has 5 independent elements so (4.20) is the  $5P$ -parameter solution.

## 5 Quantum deformations of the ADHM-solution

The covariant formulation of the ADHM multi-instanton solution in the classical twistor space was considered in Ref[13]. We shall discuss the quantum deformations of this formalism.

Let us consider the gauge group  $GL_q(N, C)$ . The ADHM-solution for the instanton number  $p$  can be connected with some  $GL_q(N+2p, C)$  matrix  $q$ -twistor function. Introduce the notation for indices of different types :  $A, B \dots = 1 \dots p$ ;  $I, K, L, M \dots = 1 \dots N+2p$  and  $i, k, l \dots = 1 \dots N$ . The ADHM Ansatz for the general self-dual  $GL_q(N, C)$  field contains the deformed twistor functions  $u_I^i(z)$  and  $\tilde{u}_i^I(z)$

$$A_k^i = du_I^i \tilde{u}_k^I, \quad u_I^i \tilde{u}_k^I = \delta_k^i \quad (5.1)$$

The commutation relations for the  $u$  and  $\tilde{u}$  twistors are

$$(R_N)_{lm}^{ik} u_I^l u_K^m = u_L^i u_M^k \mathbf{R}_{IK}^{LM} \quad (5.2)$$

$$\mathbf{R}_{ML}^{KI} \tilde{u}_i^L \tilde{u}_k^M = \tilde{u}_l^I \tilde{u}_m^K (R_N)_{ki}^{ml} \quad (5.3)$$

$$\tilde{u}_l^I (R_N)_{mk}^{li} u_K^m = u_L^i \mathbf{R}_{KM}^{IL} \tilde{u}_k^M \quad (5.4)$$

where the  $R$ -matrices for  $GL_q(N, C)$  and  $GL_q(N + 2p, C)$  are used.

Introduce also the relation for the differentials  $du$

$$\tilde{u}_i^I (R_N)_{lm}^{ik} du_K^l = du_L^k (\mathbf{R}^{-1})_{KM}^{IL} \tilde{u}_m^M \quad (5.5)$$

$$du_L^i du_M^k (\mathbf{R}^{-1})_{IK}^{LM} = -(R_N^{-1})_{lm}^{ik} du_l^I du_k^m \quad (5.6)$$

These relations are necessary for proving a validity of the gauge algebra (3.6) in the framework of the ADHM-Ansatz

$$(AR_N A)_{mn}^{ik} = du_I^i du_L^k (\mathbf{R}^{-1})_{KM}^{IL} \tilde{u}_n^M \tilde{u}_m^K = -(R_N A R_N A R_N)_{mn}^{ik} \quad (5.7)$$

Consider also the linear twistor functions  $v$  and  $\tilde{v}$

$$v_I^{A\alpha} = z_a^\alpha b_I^{aA} \quad (5.8)$$

$$\tilde{v}^{IA\alpha} = z_a^\alpha \tilde{b}^{aIA} \quad (5.9)$$

where  $b$  and  $\tilde{b}$  are the noncommutative  $q$ -instanton moduli

$$b_I^{aA} z_b^\alpha = R_{cb}^{da} z_d^\alpha b_I^{cA} \quad (5.10)$$

$$\tilde{b}^{aIA} z_b^\alpha = R_{cb}^{da} z_d^\alpha \tilde{b}^{cIA} \quad (5.11)$$

The relations between  $b$  and  $\tilde{b}$  will be defined below.

Introduce the following condition for the functions  $v$  and  $\tilde{v}$ :

$$v_I^{A\alpha} \tilde{v}^{IB\beta} = g^{AB}(z) \varepsilon^{\alpha\beta}(q) \quad (5.12)$$

where  $g(z)$  is the nondegenerate  $(p \times p)$  matrix with the central elements

$$g^{AB}(z) = q^{-2} y_{cd} b_I^{cA} \tilde{b}^{dIB} \quad (5.13)$$

The condition (5.12) is equivalent to the restriction on the elements of the  $B$ -algebra

$$[P^{(+)}]_{cd}^{ab} b_I^{cA} \tilde{b}^{dIB} = 0 \quad (5.14)$$

Write the basic commutation relations of the  $B$ -algebra

$$R_{cd}^{ab} b_I^{cA} b_K^{dB} = b_L^B b_M^A \mathbf{R}_{KI}^{ML} \quad (5.15)$$

$$\mathbf{R}_{LM}^{IK} \tilde{b}^{aLA} \tilde{b}^{bMB} = R_{cd}^{ab} \tilde{b}^{cIB} \tilde{b}^{dKA} \quad (5.16)$$

$$R_{cd}^{ab} b_I^{cA} \tilde{b}^{dKB} = \mathbf{R}_{IM}^{KL} \tilde{b}^{aMB} b_L^{bB} \quad (5.17)$$

Remark that a formal permutation of the indices  $A$  and  $B$  is commutative.

Consider the new functions

$$\tilde{v}_{A\alpha}^I = \tilde{v}^{IB\beta} g_{BA}(z) \varepsilon_{\beta\alpha}(q) \quad (5.18)$$

where we use the matrix  $g_{BA}$  inverse of the matrix (5.13)

$$g_{BA}(z) g^{AC}(z) = \delta_B^C \quad (5.19)$$

Now one can construct the full quantum  $GL_q(N+2p, C)$  matrices

$$\mathbf{U} = \begin{pmatrix} u_I^i \\ v_I^{A\alpha} \end{pmatrix}, \quad \mathbf{S}(\mathbf{U}) = \mathbf{U}^{-1} = \begin{pmatrix} \tilde{u}_i^I \\ \tilde{v}_{A\alpha}^I \end{pmatrix} \quad (5.20)$$

The standard  $GL_q(N+2p, C)$  commutation relations for these matrices contain Eqs(5.2-5.4) and the relations for  $v$  and  $\tilde{v}$  functions

$$\widetilde{\mathbf{R}} \mathbf{U} \mathbf{U}' = \mathbf{U} \mathbf{U}' \mathbf{R} \quad (5.21)$$

$$\mathbf{R} \mathbf{S}' \mathbf{S} = \mathbf{S}' \mathbf{S} \widetilde{\mathbf{R}} \quad (5.22)$$

$$\mathbf{S} \widetilde{\mathbf{R}} \mathbf{U} = \mathbf{U}' \mathbf{R} \mathbf{S}' \quad (5.23)$$

where the  $R$ -matrix for  $GL_q(N+2p, C)$  can be written in the following form

$$\widetilde{\mathbf{R}} = \begin{pmatrix} (R_N)_{mn}^{ik} & 0 & 0 & 0 \\ 0 & \lambda \delta_C^A \delta_\mu^\alpha \delta_n^k & \delta_D^A \delta_\nu^\alpha \delta_m^k & 0 \\ 0 & \delta_n^i \delta_C^B \delta_\mu^\beta & 0 & 0 \\ 0 & 0 & 0 & \delta_D^A \delta_C^B R_{\mu\nu}^{\alpha\beta} \end{pmatrix}$$

where  $\lambda = q - q^{-1}$ .

The equations (5.5,5.6) follow from are the relations

$$\mathbf{S} \widetilde{\mathbf{R}} d\mathbf{U} = d\mathbf{U}' \mathbf{R}^{-1} \mathbf{S}' \quad (5.24)$$

It should be stressed that the bicovariant differential calculus on  $GL_q(N+2p, C)$  is the basis of the deformed ADHM-construction for the group  $GL_q(N, C)$ .

Write explicitly the orthogonality and completeness conditions for the deformed ADHM-twistors:

$$u_I^i \tilde{v}^{IA\alpha} = 0 \quad (5.25)$$

$$v_I^{A\alpha} \tilde{u}_i^I = 0 \quad (5.26)$$

$$\delta_K^I = \tilde{u}_i^I u_K^i + \tilde{v}^{IA\alpha} g_{AB}(z) \varepsilon_{\alpha\beta}(q) v_K^{B\beta} \quad (5.27)$$

Now we are in a position to verify the self-duality of the connection (5.1)

$$\begin{aligned} dA_k^i - A_l^i A_k^l &= du_I^i (\tilde{u}_l^I u_M^l - \delta_M^I) d\tilde{u}_k^M = \\ &= -q^{-4} u_I^i \tilde{b}^{cIA} D_c^a g_{AB}(z) \varepsilon_{\alpha\beta}(q) dz_a^\alpha dz_b^\beta b_M^{bB} \tilde{u}_k^M \end{aligned} \quad (5.28)$$

where  $D_c^a$  is the  $GL_q(4)$  metric. This curvature contains only the self-dual  $q$ -twistor 2-form (2.21).

The real forms of the deformed ADHM-construction are based on the quantum groups  $U_q(N)$  and  $GL_q(N, R)$ .

It should be stressed that all  $R$ -matrices of our deformation scheme satisfy the Hecke relation with the common parameter  $q$ . The other possible parameters of different  $R$ -matrices are independent. The case  $q = 1$  corresponds to the unitary deformations ( $R^2 = I$ ) of the twistor space and the gauge groups. It is evident that the trivial deformation of the  $z$ -twistors is consistent with the nontrivial unitary deformation of the gauge sector and vice versa.

The author would like to thank A.T. Filippov, E.A. Ivanov, A.P.Isaev and V.I. Ogievetsky for helpful discussions and interest in this work.

I am grateful to the administration of JINR and Laboratory of Theoretical Physics for hospitality. This work was supported in part by the ISF-grant RUA000, INTAS-grant 93-127 and the contract No.40 of Uzbek Foundation of Fundamental Research .

## References

- [1] A.Connes, M.Rieffel, Contemp.Math., **62** (1987) 237
- [2] I.Ya.Aref'eva, I.V.Volovich, Mod. Phys. Lett., **A6** (1991) 893
- [3] T.Brzezinski, Sh.Majid , Commun. Math. Phys., **157** (1993) 591
- [4] M.J.Pflaum, Comm.Math.Phys., **166** (1994) 279
- [5] A.P.Isaev, Z.Popowicz , Phys.Lett., **B281** (1992) 271 ; Phys.Lett., **B307** (1993) 353
- [6] A.P.Isaev, J.Math.Phys., **35** (1995) 6784
- [7] B.M.Zupnik, JETP Lett., **61** (1995) 443; Preprints JINR E2-94-449, hep-th/9411186; E2-94-487, q-alg/9412010
- [8] V.Lakshmibai , N.Yu.Reshetikhin, C.R. Acad.Sci. Paris, **313** (1991) 121; Contemp. Math., **134** (1992) 145
- [9] S. Levendorski, Ya.Soibelman, Comm.Math.Phys., **139** (1991) 141
- [10] S.A.Merkulov, Z.Phys., **C52** (1991) 583
- [11] A.A.Belavin, A.M.Polyakov, A.S.Schwartz and Yu.S.Tyupkin, Phys.Lett., **B59** (1975) 85
- [12] M.F.Atiyah, V.G.Drinfel'd, N.J.Hitchin and Yu.I.Manin, Phys. Lett., **A65** (1978) 185  
Yu.I.Manin, Gauge Fields and Complex Geometry, Nauka Publ., Moscow, 1984
- [13] W.Siegel, Preprint ITP-SB-94-66, Stony Brook, 1994; hep-th/9412011
- [14] B.M.Zupnik, Pis'ma ZhETF, **62** (1995) 365; q-alg/9507024

- [15] N.Yu.Reshetikhin, L.A.Takhtadjan, L.D.Faddeev, *Algeb. Anal.*, **1** (1989) 178
- [16] D.D.Demidov, Yu.I.Manin, E.E.Mukhin and D.V.Zhdanovich, *Prog. Theor. Phys. Suppl.*, **102** (1990) 203
- [17] A.Sudbery, *J.Phys.*, **A23** (1990) L697
- [18] A.Schirrmacher, *Z.Phys.*, **C50** (1991) 321
- [19] B.M.Zupnik, *Teor.Mat.Phys.*, **95** (1993) 403, hep-th/9211065; *Symmetry methods in physics*, v.2, p.599, E2-94-347, (Dubna, 1994), hep-th/9307149
- [20] S.L.Woronowicz, *Comm. Math. Phys.*, **122** (1989) 125
- [21] A.Sudbery, *Phys. Lett.*, **B284** (1992) 61
- [22] P.Schupp, P.Watts, B.Zumino, *Comm. Math. Phys.*, **157** (1993) 305
- [23] A.Schirrmacher, in: *Groups and related topics*, ed. R.Gielerak et al. (Kluwer Acad. Publ. 1992) p.55
- [24] M.Chaichian, A.P.Demichev, *Phys. Lett.*, **304B** (1994) 220
- [25] V.K.Dobrev, *Phys. Lett.*, **341B** (1994) 133
- [26] G. t'Hooft, unpublished
- [27] R.Jackiw, C.Hohl, C.Rebbi, *Phys. Rev.*, **D15** (1977) 1672
- [28] E.Corrigan, D.Fairlie, *Phys.Lett.*, **67B** (1977) 69